

Spastic Colon Medication

S **pastic colon** is a common disorder of the bowel portion, which causes signs like, pain in the abdomen and irregular bowel movements. Commonly referred to as **irritable bowel syndrome** (IBS), it is associated with repeated contractions of the small and large intestinal muscles. Despite the fact that there is no cure, the condition can be combated with effective medication and dietary changes.

- You are a fan of the taste and consistency of Greek yogurt, it can be an essential part to weight management.
- Its higher protein product will keep you complete longer.
- The lower sugar is a great choice for diabetics and others looking to decrease sugar consumption.



Try Eating Regular Meals and Keeping Your Mealtimes on a Schedule

Eating on the run or missing a meal could throw off your digestive system. Take a look at your diet, and make sure you are eating the right foods. High-fiber foods, such as bran, whole grains and lots of fruits and vegetables, are good choices. It is best to avoid over-processed foods with little fiber. Also prevent large amounts of milk, cheese, meat, fatty foods and refined sugar.



ConstipationDiarrheaIrritable Bowel SyndromeIbsNauseaDiagnosis

“



Digestive System Diseases Acid reflux, hemorrhoids, irritable bowel syndrome, ulcers, etc., are the most common digestion problems. The human gastrointestinal system includes esophagus, stomach, liver, pancreas, large intestinal tract, small intestinal tract, bowel, and...

Treatment and Medication

There is no particular diagnostic method for discovering this condition. Hence diagnosis of irritable bowel syndrome is performed in the reverse way, by ruling out other health conditions that mimic **spastic colon symptoms**. For confirmation, the doctor may perform laboratory tests such as abdominal X-ray, blood test, rectal exam, colonoscopy, and urinalysis.

Enlarged Spleen

Spleen, the triangular shaped organ does an important job of producing white blood cells to fight infections. It is located in the lower left area of the chest, to the left hand side of the stomach. Damage to this vital organ due to an injury or spleen disorders makes it swollen or enlarged, leading to persistent pain under left rib cage. Mononucleosis (viral illness) and liver disorders (cirrhosis) are some of the factors that can cause spleen enlargement.

Disclaimer: Content in this document is an advertisement, therefore it may be biased and should not be considered an objective or independent review. Owner of this document may be compensated when you purchase a product by clicking a link in this document. The views and opinions expressed in this document are purely of the author. Any product claim or other representation about a product or service should be verified with the manufacturer, provider or party in question. Any health information in this document is for educational purposes only and is not intended to replace the advice of your health care provider.